arXiv:1204.2963v1 [math.CA] 13 Apr 2012

ELEMENTS OF PÓLYA-SCHUR THEORY IN FINITE DIFFERENCE SETTING

PETTER BRÄNDÉN, ILIA KRASIKOV, AND BORIS SHAPIRO

Abstract. In this note we attempt to develop an analog of Pólya-Schur theory describing the class of univariate hyperbolicity preservers in the setting of linear finite difference operators. We study the class of linear finite difference operators preserving the set of real-rooted polynomials whose mesh (i.e. the minimal distance between the roots) is at least one. In particular, finite difference versions of the classical Hermite-Poulain theorem and generalized Laguerre inequalities are obtained.

1. Introduction

Systematic study of linear operators acting on $\mathbb{R}[x]$ and sending real-rooted polynomials to real-rooted polynomials was (apparently) initiated in the 1870's by C. Hermite and later continued by E. Laguerre. Its classical period culminated in 1914 with the publication of the outstanding paper [16] where G. Pólya and I. Schur completely characterized all such linear operators acting diagonally in the standard monomial basis $(1, x, x^2, ...)$ of $\mathbb{R}[x]$. This article generated a substantial amount of related literature with important contributions of N. Obreschkov, S. Karlin, G. Csordas, T. Craven, A. Pinkus, A. Iserles, A. Bakan, and, recently by the first author together with the late J. Borcea.

Although several variations of the original set-up were considered (including complex zero decreasing sequences, real-rooted polynomials on finite intervals, stable polynomials etc.) it seems that its natural finite difference analog discussed below has so far escaped the attention of the specialists in the area. An exception is [7].

Denote by $\mathcal{HP} \subset \mathbb{R}[x]$ the set of all real-rooted (also referred to as *hyperbolic*) polynomials. A linear operator $T : \mathbb{R}[x] \to \mathbb{R}[x]$ is called a *real-rootedness preserver* or a *hyperbolicity preserver* if it preserves $\mathcal{HP}$. (We will also use the short-hand 'HPO' for such operators.) Given a real-rooted polynomial $p(x) \in \mathcal{HP}$ denote by $mesh(p)$ its mesh, i.e. the minimal distance between its roots. (If a real-rooted $p(x)$ has a multiple root then $mesh(p) = 0$.) Denote by $\mathcal{HP}_{\geq M} \subset \mathcal{HP}$ the set of all real-rooted polynomials whose mesh is at least $M > 0$. Let $\mathcal{HP}^{+}_{\geq M} \subset \mathcal{HP}_{\geq M}$ be the subset of such polynomials with only nonnegative zeros.

One of (not that many) results about linear operators not decreasing the mesh is due originally to M. Riesz and deserves to be better known, see e.g. [7], [17].

Theorem 1. *For any hyperbolic polynomial p and any real λ one has*

$$mesh(p - \lambda p') \geq mesh(p).$$

Recall that the well-known Hermite-Poulain theorem, (see [13], p. 4) claims that a finite order linear differential operator $T = a_0 + a_1 d/dx + \cdots + a_k d^k/dx^k$ with



FIGURE 1. Roots of $\Delta(p(x)) = p(x) - p(x-1)$ are the x-coordinates of the intersection points between the graphs of $p(x)$ and $p(x-1)$.

constant coefficients is a hyperbolicity preserver if and only if its symbol polynomial $Q(t) = a_0 + a_1 t + \cdots + a_k t^k$ is hyperbolic.

Thus, Theorem 1 combined with the Hermite-Poulain theorem imply the following statement.

Corollary 1. *Hyperbolicity preserving differential operator with constant coefficients does not decrease the mesh of hyperbolic polynomials.*

Our first goal is to find an analog of Corollary 1 in the finite difference context. We consider the action on $\mathbb{C}[x]$ of linear finite difference operators T with polynomial coefficients, i.e. operators of the form:

$$T(p(x)) = a_0(x)p(x) + a_1(x)p(x-1) + \cdots + a_k(x)p(x-k), \tag{1}$$

where $a_0(x), \ldots, a_k(x)$ are some fixed complex- or real-valued polynomials. Although no such T can preserve $\mathcal{HP}$, (see Lemma 7 below) it can nevertheless preserve $\mathcal{HP}_{\geq 1}$. The simplest example of such an operator is $\Delta(p(x)) = p(x) - p(x-1)$ which is a discrete analog of d/dx, see Fig. 1.

Definition 1. A linear finite difference operator (1) is called a *discrete hyperbolicity preserver* if it preserves $\mathcal{HP}_{\geq 1}$.

Obviously, the set of all discrete hyperbolicity preservers is a semigroup w.r.t. composition. We start with a finite difference analog of Theorem 1. A similar result was proved by Fisk in [7, Lemma 8.27].

Theorem 2. *For any real α, β of the same sign define the operator $T(p(x)) = \alpha p(x) - \beta p(x-1)$. Then for any hyperbolic polynomial $p(x) \in \mathcal{HP}_{\geq 1}$ one has*

$$mesh(T(p(x))) \geq mesh(p).$$

Moreover, if $\frac{\beta}{\alpha} \geq 1$ the operator T preserves the set $\mathcal{HP}^{+}_{\geq 1}$.

This statement settles Conjecture 2.19 from a recent preprint [5]. Our next result is a natural finite difference analog of the Hermite-Poulain theorem.

Theorem 3. *A linear finite difference operator T with constant coefficients of the form:*

$$T(p(x)) = a_0 p(x) + a_1 p(x-1) + \cdots + a_k p(x-k) \tag{2}$$

is a discrete hyperbolicity preserver if and only if all roots of its symbol polynomial $Q(t) = a_0 + a_1 t + \cdots + a_k t^k$ are non-negative.

As we mentioned above a famous class of hyperbolicity preservers is the class of the so-called multiplier sequences introduced by G. Pólya and I. Schur in [16]. It consists of hyperbolicity preservers acting diagonally in the standard monomial basis of $\mathbb{R}[x]$. In § 3 we present some initial definitions and results about an analog of multiplier sequences in the finite difference setting. In particular, we prove the following claim.

Theorem 4. *For any real* α, β *of the same sign define the operator* $U(p(x)) = \alpha p(x) + \beta x \Delta(p(x)) = \alpha p(x) + \beta x(p(x) - p(x-1))$. *Then* U *preserves the set* $\mathcal{HP}^{+}_{\geq 1}$.

Remark. Observe that, in general, the operator U as above is **not** mesh-increasing and, therefore, Theorem 4 is not a complete analog of Theorem 2. A simple example of this phenomenon is $U(p(x)) = p(x) + \frac{3}{4}x(p(x) - p(x-1))$, i.e. $\alpha = 1, \quad \beta = \frac{3}{4}$. When $p(x) = (x-1)(x-4)(x-7)$ then $U(p(x))$ has three positive roots which are approximately equal to $(0.433167, 3.12467, 6.36524)$ and its mesh is smaller than 3. Apparently the complete description of finite difference analogs of multiplier sequences is a difficult problem.

Hyperbolic polynomials with a restriction on their mesh play an important role in many applications. Thus, the classes $\mathcal{HP}_{\geq M}$ for $M = 1$ and 2 naturally appear in connection with discrete orthogonal polynomials, i.e., the polynomials with the orthogonality measure supported on a subset of integers. Namely, the following is known, see [12].

Theorem 5. *Let* $p(x)$ *be a discrete orthogonal polynomial. Suppose that* $p(x)$ *satisfies the difference equation*

$$p(x+1) = 2A(x)p(x) - B(x)p(x-1),$$

and there exists an open interval $I \subset \mathbb{R}$ *such that*

i) *All zeros of* p *lie in* I,
ii) $B(x) > 0$ *for* $x \in I$.

Then $mesh(p) > 1$. *If, in addition,* $A(x) > 0$ *on* I *then* $mesh(p) > 2$.

This theorem sheds some light on the concept of the discrete discriminant

$$D(p_n) = c^{2n-2} \prod_{1 \leq j < k \leq n} \left((x_j - x_k)^2 - 1 \right), \quad p_n(x) = c \prod_{j=1}^{n} (x - x_j),$$

introduced by M.E.H. Ismail (see [10, sec. 6.4] and the references therein). If p_n depends on a parameter then the orthogonality breaks down exactly when the discrete discriminant vanishes, i.e. the mesh of p_n becomes equal to one.

An important characterization of hyperbolic polynomials (or, more generally, of entire functions belonging to the Polya-Laguerre class) is given by the following set of generalized Laguerre inequalities

$$L_m[p](x) = \sum_{j=0}^{m} (-1)^{j+m} \binom{2m}{j} p^{(j)}(x) p^{(2m-j)}(x) \geq 0, \quad m = 0, 1, 2, ... \tag{3}$$

In other words, a polynomial $p(x)$ is hyperbolic if and only if (3) holds, see [6] and also [2] for a generalization.

For polynomials in $\mathcal{HP}_{\geq M}$ with $M \geq 2$ (or, more precisely, $M \geq \sqrt{4 - \frac{6}{m+2}}\,,\ \ m = 0, 1, 2, ...$), the following analog of generalized Laguerre inequalities [14], [15]

$$\sum_{j=-m}^{m} (-1)^j \frac{p(x-j)p(x+j)}{(m-j)!(m+j)!} \geq 0$$

was earlier found in [12].

In the present paper we extend the latter result by proving a new version of generalized Laguerre inequalities in the discrete setting.

Theorem 6. *Given* $0 \leq \lambda < \delta$ *let* $p(x)$ *be a hyperbolic polynomial satisfying the condition* $mesh(p) \geq \sqrt{2\lambda^2 + 2\delta^2}$. *Then one has that*

$$F_{\lambda,\delta}(p, x) = p(x-\lambda)p(x+\lambda) - p(x-\delta)p(x+\delta) \geq 0. \tag{4}$$

Moreover, the above bound on $mesh(p)$ *is sharp.*

Acknowledments. The authors are grateful to Professors O. Katkova and A. Vishnyakova of Kharkov National University for discussions of the topic. The third author is grateful to the Department of Mathematics, Brunel University for the hospitality in June 2009 when this project was initiated.

2. Proving Hermite-Poulain theorem

Lemma 7. *No finite difference operator* T *of the form* (1) *except the identity operator is hyperbolicity preserving in the classical sense.*

Proof. Notice that the main criterion of [1] implies that a linear operator $T : \mathbb{R}[x] \to \mathbb{R}[x]$ commuting with arbitrary shifts of independent variable $x \mapsto x + \gamma$ is an HPO if and only it sends each monomial $x^N,\ N = 0, 1, \dots$ to a hyperbolic polynomial. An equivalent criterion for an operator commuting with arbitrary shifts to be a HPO is that its symbol belongs to the Laguerre-Pólya class. The symbol of $T(p) = a_0p(x) + a_1p(x-1) + ... + a_kp(x-k)$ equals $S(t) = a_0 + a_1e^{-t} + ... + a_ke^{-kt}$. Such a function can not be in the Laguerre-Pólya class unless $k = 0$. □

Below we present two different proofs of Theorem 2 based on different ideas and illustrating different aspects of this result.

First proof of Theorem 2. If suffices to consider the family of operators $T_\lambda(p) = p(x) - \lambda p(x-1)$ with $\lambda > 0$ since the degenerate cases $\alpha = 0$ or $\beta = 0$ are obvious. Set $q = T_\lambda(p)$ where $p \in \mathcal{HP}_{\geq 1}$ and $\deg p = n$. The next statement is obvious.

Lemma 8. *For* $p \in \mathcal{HP}_{\geq 1}$ *and any real* λ *all roots of* q *are real and interlacing the roots of* p.

Now let $\alpha_1 < \alpha_2 < ... < \alpha_n$ be the roots of p and $\beta_1 < \beta_2 < ... < \beta_n$ be the roots of q. (Here we assume that $\lambda \neq 1$. The latter case follows by continuity.) Notice that by Lemma 8 the roots $\alpha_1 < \alpha_2 < ... < \alpha_n$ and $\beta_1 < \beta_2 < ... < \beta_n$ interlace. Set

$$t(x) = \frac{p(x-1)}{p(x)} = 1 + \sum_{j=1}^{n} \frac{c_j}{x - \alpha_j}.$$

One can easily see that all $c_j \leq 0$. Therefore, $t(x)$ is increasing between any two consecutive roots of $p(x)$. Assume first that $deg\, p = 2$. W.l.o.g. we can take $p = x(x-\alpha)$. If $\beta_1 < \beta_2$ are the roots of q then for $\lambda > 0, \lambda \neq 1$, one can easily check that

$$(mesh(q))^2 - (mesh(p))^2 = (\beta_2 - \beta_1)^2 - \alpha^2 = \frac{4\lambda}{(1-\lambda)^2} \geq 0.$$

Considering the general case $deg\, p \geq 3$, we observe first that

$$\frac{q(x)}{p(x)} = 1 - \lambda \prod_{j=1}^{n} \left(1 - \frac{1}{x-\alpha_j}\right). \tag{5}$$

Notice that the product in the r.h.s. of (5) may contain just one non-positive factor $\left(1 - \frac{1}{x-\alpha_j}\right)$ exactly when $\alpha_j < x \leq \alpha_j + 1$. However, if $q(x) = 0$ then all these factors must be strictly positive since if such a negative factor exists the r.h.s. of (5) is strictly positive. Thus, if $q(x) = 0$ and $x > \alpha_j$ then $x > \alpha_j + 1$. Therefore, for any two consecutive zeros β_i and β_{i+1} of q there exists $1 \leq l \leq n$ such that

$$\alpha_{l-1} + 1 < \beta_i < \alpha_l < \alpha_l + 1 < \beta_{i+1} < \alpha_{l+1}, \tag{6}$$

where we set $\alpha_0 = -\infty$, $\alpha_{n+1} = \infty$. (One can say more about the relation between i and l but we do not need this information in our argument.)

Let us show that $mesh(q) \geq mesh(p)$. Assume that there exist two consecutive zeros $\beta_i < \beta_{i+1}$ of q satisfying $\beta_{i+1} - \beta_i < mesh(p)$. To get a contradiction it is enough to show that

$$F = \frac{1}{\lambda}\left(\frac{q(\beta_{i+1})}{p(\beta_{i+1})} - \frac{q(\beta_i)}{p(\beta_i)}\right) \neq 0.$$

Then β_i and β_{i+1} cannot be consecutive zeros of q. We have

$$F = \prod_{j=1}^{n}\left(1 - \frac{1}{\beta_i - \alpha_j}\right) - \prod_{j=1}^{n}\left(1 - \frac{1}{\beta_{i+1} - \alpha_j}\right) =$$

$$\left(1 - \frac{1}{\beta_i - \alpha_n}\right)\prod_{j=1}^{n-1}\left(1 - \frac{1}{\beta_i - \alpha_j}\right) - \left(1 - \frac{1}{\beta_{i+1} - \alpha_1}\right)\prod_{j=2}^{n}\left(1 - \frac{1}{\beta_{i+1} - \alpha_j}\right) =$$

$$\left(1 - \frac{1}{\beta_i - \alpha_n}\right)\prod_{j=1}^{n-1}\left(1 - \frac{1}{\beta_i - \alpha_j}\right) - \left(1 - \frac{1}{\beta_{i+1} - \alpha_1}\right)\prod_{j=1}^{n-1}\left(1 - \frac{1}{\beta_{i+1} - \alpha_{j+1}}\right) :=$$

$$:= v_0 \prod_{j=1}^{n-1} v_j - u_0 \prod_{j=1}^{n-1} u_j,$$

where all v_j and u_j are positive. Finally, observe that $v_0 > 1$, $0 < u_0 < 1$, and $v_j, u_j > 0$ for $j \neq l-1$, and moreover,

$$v_j - u_j = \frac{1}{\beta_{i+1} - \alpha_{j+1}} - \frac{1}{\beta_i - \alpha_j} = \frac{(\alpha_{j+1} - \alpha_j) - (\beta_{i+1} - \beta_i)}{(\beta_i - \alpha_j)(\beta_{i+1} - \alpha_{j+1})} > 0.$$

Hence, $v_0 \prod_{j=1}^{n-1} v_j > u_0 \prod_{j=1}^{n-1} u_j$, and the result follows.

Let us finally prove that for $\lambda \geq 1$ the operator T_λ preserves the set $\mathcal{HP}^+_{\geq 1}$. Notice that the equation $q(x) = p(x) - \lambda p(x-1) = 0$ is equivalent to the equation $t(x) = \frac{1}{\lambda}$ where $t(x)$ is defined above. Denote by $B_0, B_1, ..., B_n$ the consecutive (monotone increasing) branches of $t(x)$ between its poles, i.e. the roots of $p(x)$.

In other words, B_0 is the restriction of $t(x)$ to $(-\infty,\alpha_1)$, B_1 is the restriction to (α_1,α_2), ... etc. Since all the zeros of $p(x)$ are non-negative, the line $y=\frac{1}{\lambda}$ for $\lambda>1$ intersects all the branches $B_1,...,B_n$ of $t(x)$ for some positive values of x, and does not intersect B_0 as it lies above the level $y=1$. Hence, all the zeros of $q(x)$ are positive. Finally, if $\lambda=1$ then one root escapes to ∞ and the line $y=1$ intersects the branches $B_1,...,B_{n-1}$. $\square$

Alternative proof of Theorem 2. A pair of hyperbolic polynomials (p,q) are in *proper position*, written $p\ll q$ if their zeros interlace and $p(x)q'(x)-p'(x)q(x)\geq 0$ for all $x\in\mathbb{R}$. Note that if the zeros of two hyperbolic polynomials p and q interlace then either $p\ll q$ or $q\ll p$. By convention $0\ll p$ and $p\ll 0$ for any hyperbolic polynomial p. The next lemma is well known, part (a) is almost obvious from the interlacing property and part (b) is often referred to as Obreschkoff's theorem, see [13].

Lemma 9. (a) Let p be a hyperbolic polynomial. Then the sets

$$\{q\in\mathbb{R}[x]:q\ll p\}\quad\text{and}\quad\{q\in\mathbb{R}[x]:p\ll q\}$$

are convex cones.
(b) If $p\ll q$ then $p\ll q+\alpha p$ and $p+\alpha q\ll q$ for all $\alpha\in\mathbb{R}$.

Let $T(p)(x)=p(x)-\lambda p(x-\alpha)$ where $\alpha,\lambda\geq 0$. We want to prove that $T:\mathcal{HP}_{\geq\beta}\to\mathcal{HP}_{\geq\beta}$ for all $\beta\geq\alpha$. First we prove it for $\beta=\alpha$. Note that $p\in\mathcal{HP}_{\geq\alpha}$ if and only if $p(x)\ll p(x-\alpha)$. Lemma 9 (b) implies $T(p)\ll p(x)$ and $T(p)\ll p(x-\alpha)$, which is easily seen to imply $T(p)\in\mathcal{HP}_{\geq\alpha}$. Next we prove that if $p,q\in\mathcal{HP}_{\geq\alpha}$ satisfy $p\ll q$, then $T(p)\ll T(q)$. This will prove Theorem 2, since if $p\in\mathcal{HP}_{\geq\beta}\subseteq\mathcal{HP}_{\geq\alpha}$, then $p(x)\ll p(x-\beta)$ and thus $T(p)(x)\ll T(p)(x-\beta)$ which is equivalent to $T(p)\in\mathcal{HP}_{\geq\beta}$. To prove that T preserves proper position it is enough to prove that $T((x-a)p)\ll T((x-b)p)$ whenever $a\leq b$ and $p\in\mathcal{HP}_{\geq\alpha}$. Since $T((x-b)p)=T((x-a)p)-(b-a)T(p)$ it is, by Lemma 9 (a), enough to prove that $T(p)\ll T(xp)$ for all $p\in\mathcal{HP}_{\geq\alpha}$. Now

$$T(xp)=(x-\alpha)T(p)+\alpha p.$$

Since $T(p)\ll p$ and $T(p)\ll(x-\alpha)T(p)$, Lemma 9 b implies $T(p)\ll(x-\alpha)T(p)+\alpha p$ as desired. $\square$

Corollary 2. *For any $\lambda\geq 1$ the operator $xT_\lambda(p)=x(p(x)-\lambda p(x-1))$ preserves the set $\mathcal{HP}^{+}_{\geq 1}$.*

Proof. By Theorem 2 it is enough to check that

$$q(x)=xT_\lambda(p)=x\left(p(x)-\lambda p(x-1)\right)\neq 0$$

for $x\leq 1$. Indeed, the condition $q(x)=0$ can be rewritten as $t(x)=\frac{1}{\lambda}$. For $\lambda\geq 1$ the function $y=\frac{1}{\lambda}$ intersects all the branches $B_1,...,B_{n-1}$. It also intersects B_n if $\lambda>1$. On the other hand, it does not intersect B_0 since $t(x)$ is a continuous increasing function for $x<0$ with $\lim_{x\to-\infty}t(x)=1$. Hence the least zero β of q is in the interval $[\alpha_1,\alpha_2]$, and, therefore, $\beta>\alpha_1+1>1$, by (6). $\square$

Proof of Theorem 3. Theorem 2 implies that if the symbol polynomial $Q(t)=a_0+a_1t+...+a_kt^k$ has all real and non-negative roots then the finite difference operator $T(p(x))=a_0p(x)+a_1p(x-1)+...+a_kp(x-k)$ is a discrete hyperbolicity preserver.

We need to prove the necessity of the latter condition. Consider the action of T on the Pochhammer polynomial $Po_N(x)$ defined by

$$Po_0(x) = 1, \quad Po_m(x) = x(x-1)...(x-m+1),\ m = 1, 2, ... \tag{7}$$

Assuming that $N \geq k$ we get

$$T(Po_N(x)) = (x-k)...(x-N+1)R_N(x),$$

where $R_N(x)$ is a hyperbolic polynomial of degree k. Observe that

$$\lim_{N\to+\infty} \frac{R_N(xN)}{N^k} = x^k Q\left(\frac{x}{x-1}\right),$$

where $Q(t)$ is the above symbol polynomial. Hence if T is a discrete hyperbolicity preserver then $Q(t)$ has to be hyperbolic. We need to show that its zeros are non-negative. Suppose that $Q(y) = 0$ for $y < 0$. Then the assumption $y = \frac{x}{x-1}$ implies that $0 < x < 1$. Hence, for N large enough $R_N(x)$ will have a zero in the interval $[k, N_1]$. This means that the mesh of $T(Po_N(x)) = (x-k)...(x-N+1)R_N(x)$ is strictly smaller than 1 which contradicts to our assumption. Hence all zeros of $Q(t)$ should be non-negative. □

3. On finite difference analog of multiplier sequences

Next let us recall the classical notion of a multiplier sequence due to Pólya and Schur, [16] and introduce its finite difference analog.

Definition 3. Given a sequence $\mathcal{A} = \{\alpha_n\}_{n=0}^{\infty}$ of real or complex numbers we denote by $T_{\mathcal{A}}$ the linear operator

$$T_{\mathcal{A}}(x^i) = \alpha_i x^i$$

acting diagonally in the monomial basic of $\mathbb{C}[x]$. We refer to $T_{\mathcal{A}}$ as the *diagonal operator corresponding to the sequence* $\mathcal{A}$. We call a sequence $\mathcal{A} = \{\alpha_n\}_{n=0}^{\infty}$ of real numbers a *multiplier sequence of the 1st kind*, if its diagonal operator $T_{\mathcal{A}}$ preserves $\mathcal{HP}$, i.e. sends an arbitrary hyperbolic polynomial to a hyperbolic polynomial. The above sequence $\mathcal{A}$ is called a *multiplier sequence of the 2nd kind*, if the above $T_{\mathcal{A}}$ sends an arbitrary hyperbolic polynomial whose roots are all of the same sign to a hyperbolic polynomial.

In [16] the classical criteria of being a multiplier sequence of the 1st and the 2nd kind were presented.

Definition 4. Consider a diagonal (in the monomial basis) operator $T : \mathbb{R}_n[x] \to \mathbb{R}_n[x]$ acting by multiplication of x^j by $\gamma_j,\ j = 0, \ldots, n$. A diagonal hyperbolicity preserver T will be referred to as a *multiplier sequence of length* $n+1$ or simply a *finite multiplier sequence.* Denoted by $M_n \subset \mathbb{R}^{n+1}$ the set of all finite multiplier sequences of length $n+1$.

A characterization of finite multiplier sequence similar to that of [16] can be found in [4], Theorem 3.7, see also [3], Theorem 3.1. One can identify the semigroup of all finite multiplier sequences of length $n+1$ whose polynomials $Q_T(t)$ have all positive roots with the set of such polynomials. The usual multiplication of diagonal matrices then corresponds to the so-called Schur-Szegö multiplication of polynomials, see e.g. [18]. Namely, the Schur-Szegö product $P * Q$ of two polynomials $P(x) = \sum_{j=0}^n a_j \binom{n}{j} x^j$ and $Q(x) = \sum_{j=0}^n b_j \binom{n}{j} x^j$ equals

$$P * Q = \sum_{j=0}^{n} a_j b_j \binom{n}{j} x^j.$$

Apparently there were no previously known results relating multiplier sequences and mesh. The next two theorems were recently proven in [11].

Theorem 10. *The multiplier sequence defined by the differential operator* $T(p(x)) = \lambda p + xp'$, $\lambda > 0$ *has the property*

$$\mathrm{lmesh(T(p))} > \mathrm{lmesh(p)},$$

where the logarithmic mesh of a polynomial $p(x)$ *with all positive roots* $0 < x_1 \leq x_2 \leq ... \leq x_n$ *is given by*

$$\mathrm{lmesh(p)} = \min_{\mathrm{j=1,\dots,n-1}} \frac{\mathrm{x_{j+1}}}{\mathrm{x_j}}.$$

Theorem 11. *Given two polynomials* P *and* Q *of degree* n *with all positive roots one has*

$$\mathrm{lmesh(P * Q)} \geq \mathrm{max(lmesh(P), lmesh(Q))}.$$

Let us now present a finite difference analog of multiplier sequences. A natural analog of monomials in the finite difference setting are the Pochhammer polynomials $\{Po_i(x)\}$, $i = 0, 1, ...$ given by (7). In particular, a finite difference analog of the Euler operator $x\frac{d}{dx}$ given by $x\Delta = x(p(x) - p(x-1))$ acts diagonally in this basis, namely, $x\Delta(Po_i(x)) = iP_i(x)$. This observation motivates the following notion.

Definition 5. A finite difference operator T as in (1) is called *diagonal* if it acts diagonally in the Pochhammer basis $\{Po_i(x)\}$. We say that a diagonal finite difference operator T is a *discrete multiplier sequence* if it preserves $\mathcal{HP}^+_{\geq 1}$.

Let us now prove Theorem 4. To do this it suffices to consider the operator:

$$W_\lambda(p) = p(x) + \lambda x \Delta(p(x)) = p(x) + \lambda x \left(p(x) - p(x-1)\right).$$

Proposition 12. *For any* $\lambda \geq 0$ *one has*

$$W_\lambda(p) : \mathcal{HP}^+_{\geq 1} \to \mathcal{HP}^+_{\geq 1}.$$

Proof. We first show that for any $p \in \mathcal{HP}^+_{\geq 1}$ with $\deg p(x) = n$ the polynomial $q(x) = W_\lambda(p) = p(x) + \lambda x \left(p(x) - p(x-1)\right)$ has for any $\lambda > 0$ n non-negative roots, see Fig. 2.

We consider the case when all zeros of $p(x)$ are strictly positive. (The remaining case follows by continuity.) In fact, we can prove that if $p \in \mathcal{HP}_{\geq 1}$ and $\lambda > 0$ then $q(x)$ is hyperbolic. Consider the quotient

$$\Psi(x) = \frac{q(x)}{\lambda x p(x)} = 1 + \frac{1}{\lambda x} - t(x)$$

whose zeros coincide with that of $q(x)$. Thus, roots of $q(x)$ solve the equation $t(x) = 1 + \frac{1}{\lambda x}$. Notice that the graph of $t(x)$ considered as a function on $\mathbb{R}P^1$ has no critical points and makes exactly n turns around the image $\mathbb{R}P^1$ in the counterclockwise direction while the graph of $1 + \frac{1}{\lambda x}$ for $\lambda > 0$ makes one turn in the clockwise direction and they meet on $\mathbb{R}P^1$ exactly n times.



FIGURE 2. Intersection of the graph of $t(x) = \frac{p(x-1)}{p(x)}$ for $p(x) = (x-3)(x-6)(x-9)$ with the graph of $1 + \frac{2}{x}$.

We now check that if $p \in \mathcal{HP}^+_{\geq 1}$ then $W_\lambda(p) \in \mathcal{HP}^+_{\geq 1}$. We have to check that the graphs of t and $1 + \frac{1}{\lambda x}$ do not intersect for negative values of x. This is obvious as the negative branch of the hyperbola $y = 1 + \frac{1}{\lambda x}$ does not exceed 1, whereas $t(x) > 1$ for $x < 0$.

Let us finally show that if $p \in \mathcal{HP}^+_{\geq 1}$ then the minimal distance between the roots of $q(x)$ is at least 1. Analogously to (5) one gets

$$\Psi(x) = \frac{q(x)}{\lambda x p(x)} = 1 + \frac{1}{\lambda x} - t(x) = 1 + \frac{1}{\lambda x} - \prod_{j=1}^{n} \left(1 - \frac{1}{x - \alpha_j}\right).$$

Again, since $p \in \mathcal{HP}_{\geq 1}$ the product in the right-most part of the latter formula might have at most one negative factor. Moreover, for $\lambda > 0$ no such factor can be negative at any root of $\Psi(x)$ since otherwise the right-hand side is obviously positive. Therefore, the inequality (6) is valid for the roots of $q(x)$ implying that the mesh of $q(x)$ is at least 1. □

4. PROVING DISCRETE ANALOG OF GENERALIZED LAGUERRE INEQUALITIES

We start with the following remark. Let $p(x) = \prod_{j=1}^{k}(x - x_j)$ be a hyperbolic polynomial. Consider a quadratic form

$$\mathcal{L}_m(p, x) = \sum_{i=0}^{m} a_i p^{(i)}(x) p^{(2m-i)}(x),\ a_i \in \mathbb{R}.$$

One can easily see that if $\mathcal{L}_m$ attains nonnegative values on all hyperbolic polynomials then $a_m > 0$. Indeed, choose $p(x) = h_k(x)$, where $h_k(x) = 2^k k! H_k(x)$, is a normalized Hermite polynomial. Then $h_k'(x) = h_{k-1}(x)$ and assuming that $a_m \leq 0$ we get

$$\int_{\infty}^{\infty} \mathcal{L}_m(h_k, x) e^{-x^2} dx \leq 0.$$

Thus, $\mathcal{L}_m(h_k, x) < 0$ for some x.

Theorem 6 whose proof is given below shows that the situation for polynomials with restrictions on mesh is quite different.

Proof of Theorem 6. We use induction on the degree of p. One has

$$F_{\lambda,\delta}(1, x) = 0,\ \ F_{\lambda,\delta}(x, x) = \delta^2 - \lambda^2,$$

$$F_{\lambda,\delta}(x^2-\frac{M^2}{4},x)=\frac{\delta^2-\lambda^2}{2}\,(4x^2+M^2-2\lambda^2-2\delta^2).$$

Hence Theorem 6 holds for polynomials of degree less than 3.

Consider the general case. Without loss of generality it suffices to show that $F_\lambda(p,0)\geq 0$. Assuming that (4) holds for a polynomial $p(x)$ of degree $k-1\geq 2$, consider the polynomial $q(x)=(x-x_0)p(x)$.

Notice that by the assumption on the mesh we can choose x_0 far enough from the origin. Namely, we can assume that

$$|x_0|\geq\frac{3}{2}\sqrt{2\lambda^2+2\delta^2}\,.$$

Then one has

$$F_{\lambda,\delta}(q,x)-\left((x-x_0)^2-c\right)F_{\lambda,\delta}(p,x)=$$
$$=(c-\lambda^2)p(x-\lambda)p(x+\lambda)+(\delta^2-c)p(x-\delta)p(x+\delta).$$

Choosing $c=\lambda^2$ or $c=\delta^2$, by $x_0^2>\delta^2$, we conclude that $F_{\lambda,\delta}(q,x)\geq 0$, if at least one of the expressions $p(x-\lambda)p(x+\lambda)$, $p(x-\delta)p(x+\delta)$, is nonnegative.

We need to consider the remaining case when $p(x-\lambda)p(x+\lambda)<0$, $p(x-\delta)p(x+\delta)<0$. This situation occurs only if there is a zero x_i of $p(x)$ such that $-\lambda<x_i<\lambda$. Moreover, the lower bound on the mesh implies that x_i is the only zero in the interval $[-\delta,\delta]$. Set

$$q(x)=\prod_{j=1}^{k}(x-x_j),\ \ x_1<x_2<...<x_i<...<x_k.$$

It suffices to show that in this case one has $p(-\lambda)p(\lambda)>p(-\delta)p(\delta)$, i.e.

$$\prod_{j=1}^{k}(x_j^2-\lambda^2)>\prod_{j=1}^{k}\left(x_j^2-\delta^2\right),$$

or, equivalently,

$$G(x_i)=\frac{\lambda^2-x_i^2}{\delta^2-x_i^2}\,AB\leq 1,$$

where

$$A=\prod_{j=1}^{i-1}\frac{x_j^2-\lambda^2}{x_j^2-\delta^2}\,,\ \ B=\prod_{j=i+1}^{k}\frac{x_j^2-\lambda^2}{x_j^2-\delta^2}\,.$$

Setting $\mu=\sqrt{2\lambda^2+2\delta^2}$ we get

$$A=\prod_{j=1}^{i-1}\left(1-\frac{\delta^2-\lambda^2}{\delta^2-x_j^2}\right)\leq\prod_{j=1}^{\infty}\left(1-\frac{\delta^2-\lambda^2}{\delta^2-(x_i-j\mu)^2}\right)=\frac{\Gamma\left(1-\frac{\delta+x_i}{\mu}\right)\Gamma\left(1+\frac{\delta-x_i}{\mu}\right)}{\Gamma\left(1-\frac{\lambda+x_i}{\mu}\right)\Gamma\left(1+\frac{\lambda-x_i}{\mu}\right)}\,.$$

Similarly,

$$B\leq\prod_{j=1}^{\infty}\left(1+\frac{\delta^2-\lambda^2}{(x_i+j\mu)^2-\delta^2}\right)=\frac{\Gamma\left(1-\frac{\delta-x_i}{\mu}\right)\Gamma\left(1+\frac{\delta+x_i}{\mu}\right)}{\Gamma\left(1-\frac{\lambda-x_i}{\mu}\right)\Gamma\left(1+\frac{\lambda+x_i}{\mu}\right)}\,.$$

Finally, combining these estimates and simplifying we get

$$G(x_i)\leq\frac{\sin\frac{\pi(\lambda-x_i)}{\mu}\,\sin\frac{\pi(\lambda+x_i)}{\mu}}{\sin\frac{\pi(\delta-x_i)}{\mu}\,\sin\frac{\pi(\delta+x_i)}{\mu}}\,,$$

and

$$\pi^{-1}\sin^2\frac{\pi(\delta-x_i)}{\mu}\sin^2\frac{\pi(\delta+x_i)}{\mu}G'(x_i)=\sin\frac{\pi}{\mu}\ \sin\frac{\pi(\delta^2-\lambda^2)}{\mu}\ \sin\frac{2\pi x_i}{\mu}.$$

Hence, the maximum of G is attained at $x_i=0$ and equals

$$G(0)=\frac{\sin^2\frac{\lambda\pi}{\mu}}{\sin^2\frac{\delta\pi}{\mu}}<1.$$

This proves (4).

Let us show that the bound on the mesh is sharp. We exploit the fact that the lower bound is sharp for quadratic polynomials. Let $f(x)=p(x)q(x),\ \ mesh(f)=M$, where $p(x)=x(x-M)$, and $q(x)=\prod_{i=1}^n(x-x_i)$ is a polynomial with sufficiently large zeros, such that

$$\frac{q(x-\delta)q(x+\delta)}{q(x-\lambda)q(x-\lambda)}=\prod_{j=0}^{n}\left(1-\frac{\delta^2-\lambda^2}{(x-x_i)^2-\lambda^2}\right)=1-\epsilon$$

for $x=M/2$. We can write

$$\frac{F_\lambda(f,x)}{q(x-\lambda)q(x+\lambda)}=F_\lambda(p,x)+p(x-\delta)p(x+\delta)\left(1-\frac{q(x-\delta)q(x+\delta)}{q(x-\lambda)q(x+\lambda)}\right).$$

Choosing $x=M/2$ we see that the denominator $q(\frac{M}{2}-\lambda)q(\frac{M}{2}+\lambda)$ of the left-hand side is positive, whereas the right-hand side is equal to

$$\frac{\delta^2-\lambda^2}{2}\left(M^2-2\lambda^2-2\delta^2\right)+\frac{\left(4\delta^2-M^2\right)^2}{16}\epsilon.$$

Since ϵ can be made arbitrarily small we obtain that

$$F_\lambda\left(f,\frac{M}{2}\right)<0$$

if $M<\sqrt{2(\lambda^2+\delta^2)}$. This completes the proof. □

Combining Theorem 6 with Theorem 2 one can obtain higher order difference inequalities. To avoid cumbersome expressions we only consider the simplest case corresponding to $\lambda=0,\ \delta=1$.

Corollary 3. *Let* $p(x)$ *be a hyperbolic polynomial of degree at least 2 with* $mesh(p)\geq\sqrt{5}$, *then*

$$3p^2(x)p^2(x+1)-4p(x-1)p^3(x+1)-4p^3(x)p(x+2)+$$
$$+6p(x-1)p(x)p(x+1)p(x+2)-p^2(x-1)p^2(x+2)\geq 0.$$

Proof. For $mesh(p)\geq\sqrt{2}$, we have

$$F_{0,1}(p,x)=p(x)^2-p(x-1)p(x+1)\geq 0.$$

Consider the polynomial $q(x)=p(x+1)-cp(x),\ \ c\geq 0$. We get

$$F_{0,1}(q,x)=c^2F_{0,1}(p,x)-cF_{\frac{1}{2},1}(p,x+\frac{1}{2})+F_{0,1}(p,x+1)\geq 0,$$

since $mesh(q)\geq mesh(p)$ by Theorem 2.

The minimum of $F_{0,1}(q,x)$ w.r.t. c is attained at

$$c = \frac{F_{\frac{1}{2},1}(p, x+\frac{1}{2})}{2F_{0,1}(p,x)},$$

where $c > 0$ if $mesh(p) \geq \sqrt{5}$, and is equal to

$$3p^2(x)p^2(x+1) - 4p(x-1)p^3(x+1) - 4p^3(x)p(x+2)+$$
$$+6p(x-1)p(x)p(x+1)p(x+2) - p^2(x-1)p^2(x+2).$$

□

5. Final remarks

1. Recall that the Hermite-Poulain theorem has a version in finite degrees (which we could not find in the literature). Namely, the following statement holds.

Proposition 13. *A differential operator* $T = a_0 + a_1\frac{d}{dx} + ... + a_k\frac{d^k}{dx^k}$, $a_k \neq 0$ *with constant coefficients preserves the set of hyperbolic polynomial of degree at most* m *if and only if the polynomial* $T(x^m)$ *is hyperbolic.*

The role of monomials x^m in the finite difference setting is often played by the Pochhammer polynomials (7). In particular, Proposition 13 might have the following conjectural analog in the finite difference setting.

Conjecture 1. *A difference operator* $T(p(x)) = a_0p(x)+a_1p(x-1)+...+a_kp(x-k)$ *with constant coefficients preserves the set of hyperbolic polynomial of degree at most* m *whose mesh is at least one if and only if the polynomial* $T(Po_m(x))$ *is hyperbolic and has mesh at least one.*

There is an alternative formulation of Conjecture 1 which is maybe more attractive. Let $\nabla p(x) = p(x+1) - p(x)$ be the forward difference operator, and consider the following product on the space of polynomials of degree at most d:

$$(p \bullet q)(x) = \sum_{k=0}^{d} (\nabla^k p)(0)(\nabla^{d-k})q(x).$$

Then Conjecture 1 is equivalent to

Conjecture 2. *If* p *and* q *are hyperbolic polynomials of degree at most* d *and of mesh* ≥ 1*, then so is* $(p \bullet q)$*.*

2. Problem 1. What is an analog of the famous characterization of multiplier sequence obtained in [16] in the finite difference setting?

It seems that the logarithmic mesh is more appropriate for multiplier sequence, see Theorems 10 and 11. Therefore, it is not clear how complicated an answer to the latter question might be.

3. Extending Theorem 6 we suggest the following conjecture.

Conjecture 3. *Let* $p(x)$ *be a hyperbolic polynomial with* $mesh(p) \geq 2(\lambda + \delta + \gamma)$, *where* λ, δ, γ *are nonnegative and* $\lambda \leq \delta$, *then*

$$F_{\lambda+\gamma,\delta+\gamma}(p,x) \geq F_{\lambda,\delta}(p,x). \tag{8}$$

Let us notice a special case of the above conjecture corresponding to $\lambda = 0,\ \delta = 1,\ \gamma = 1:$

$$p^2(x) - p(x-1)p(x+1) \leq p(x-1)p(x+1) - p(x-2)p(x+2), \quad mesh(p) \geq 4.$$

Let us also notice that while the extremal polynomial for (4) are just quadratic polynomials, it might happen that for (8) they are "polynomials" of an infinite degree with equi-spaced zeros, i.e. $\sin \frac{\pi x}{M}$, in which case we get

$$F_{\lambda+\gamma,\delta+\gamma}(\sin \frac{\pi x}{M}, x) - F_{\lambda,\delta}(\sin \frac{\pi x}{M}, x) = 2\cos \frac{(\lambda+\delta+\gamma)\pi}{M}\ \sin \frac{\gamma\pi}{M}\ \sin \frac{(\delta-\lambda)\pi}{M}\ .$$

References

[1] J. Borcea, P. Brändén, *Pólya-Schur master theorems for circular domains and their boundaries.* Ann. of Math. (2) 170 (2009), no. 1, 465–492.

[2] D.A. Cardon, *Extended Laguerre inequalities and a criterion for real zeros*, Progress in Analysis and its Applications, Proceedings of the 7th International Isacc Conference, (2009), 143-149.

[3] T. Craven, G. Csordas, *Problems and theorems in the theory of multiplier sequences.* Serdica Math. J. **22** (1996), 515–524.

[4] T. Craven and G. Csordas, *Multiplier sequences for fields.* Illinois J. Math. 21 (1977), no. 4, 801–817.

[5] G. Csordas, M. Chasse, *Discrete analogues of the Laguerre inequalities and a conjecture of I. Krasikov*, arXiv:1005.5186v2.

[6] G. Csordas, R.S. Varga, *Necessary and sufficient conditions and the Riemann hypothesis*, Adv. in Appl. Math. 11 (1990), 328-357.

[7] S. Fisk *Polynomials, roots, and interlacing*, arXiv:math/0612833.

[8] W.H. Foster and I. Krasikov, *Inequalities for real-root polynomials and entire functions,* Adv. Appl. Math, 29, (2002) 102-114.

[9] A. O. Gelfond, Calculus of finite differences. Translated from the Russian. International Monographs on Advanced Mathematics and Physics. Hindustan Publishing Corp., Delhi, 1971. vi+451 pp.

[10] M.E.H. Ismail, Classical and quantum orthogonal polynomials in one variable. Encyclopedia of Mathematics and its Applications, 98. Cambridge University Press, Cambridge, 2005, xviii+706 pp.

[11] O. Katkova, B. Shapiro, and A. Vishnyakova, *Multiplier sequences and logarithmic mesh.* C. R. Math. Acad. Sci. Paris 349 (2011), no. 1-2, 35–38.

[12] I. Krasikov, *Discrete analogues of the Laguerre inequality.* Analysis and Applications, 1 (2003), 189–198.

[13] N. Obreschkov, Verteilung und Berechnung der Nullstellen reeller Polynome. (German) VEB Deutscher Verlag der Wissenschaften, Berlin 1963, viii+298 pp.

[14] M. L. Patrick, *Some inequalities concerning Jacobi Polynomials.* SIAM J. Math. Anal., 2 (1971) 213-220.

[15] M. L. Patrick, *Extensions of inequalities of the Laguerre and Turn type.* Pacific J. Math. 44 (1973), 675–682.

[16] G. Pólya and J. Schur, *Über zwei Arten von Faktorenfolgen in der Theorie der algebraische Gleichungen.* J. Reine und Angew. Math., v. 144 (1914), 89–113.

[17] A. Stoyanoff, *Sur un theoreme de M Marcel Riesz.* Nouvelles Annales de Mathematique 1 (1926), 97–99.

[18] G. Szegö, *Bemerkungen zu einem Satz von J. H. Grace über die Wurzeln algebraischer Gleichungen.* Math.Z., (2) vol. 13, (1922), 28–55.

Department of Mathematics, Royal Institute of Technology, SE-100 44 Stockholm, Sweden
E-mail address: `pbranden@kth.se`

Department of Mathematical Sciences, Brunel University, Uxbridge UB8 3PH United Kingdom
E-mail address: `mastiik@brunel.ac.uk`

Department of Mathematics, Stockholm University, S-10691, Stockholm, Sweden
E-mail address: `shapiro@math.su.se`